

FOURTH QUARK-LEPTON GENERATION AND PRECISION MEASUREMENTS

M.I. VYSOTSKY

ITEP, Moscow, Russia

Precise measurements of Z -boson parameters and W -boson and t -quark masses put strong constraints on non $SU(2) \times U(1)$ singlet New Physics. We demonstrate that one extra generation passes electroweak constraints even when all new particle masses are well above their direct mass bounds. My talk is based on a recent paper¹.

1 Decoupling and nondecoupling

Let us start from Quantum Electrodynamics and consider heavy t -quark contributions to physical observables. Contribution of a diagram with top quark propagating in a loop to fine structure constant α is divergent and not extractable from data. t -quark contribution to muon anomalous magnetic moment is suppressed as $\alpha^2(m_\mu/m_t)^2$, so it decouples.

What about electroweak theory? t -quark contribute into $K - \bar{K}$, $B - \bar{B}$ transition amplitudes $\sim f^4/m_t^2 = G_F^2 m_t^2$, where f is Yukawa coupling of top with higgs boson, $f \sim m_t/\eta$, so it does not decouple. Even if new generations mixing with light generations is small (vanishing contributions to $K - \bar{K}$, $B - \bar{B}$ mixing) they contribute to Z and W polarization operators. Resulting contributions to physical observables are finite and do not decouple. This is precisely the reason why considerable part of phase space (masses of new quarks and leptons) is excluded by precision data.

2 SM fit

The precision data fit within Standard Model with three generations performed by LEPTOP code is presented in¹. The data set contains Z -boson decay parameters, W -boson and t -quark masses and the value of running fine structure constant at $q^2 = m_Z^2$.

The quality of the fit is characterized by $\chi^2/n_{d.o.f.} = 18/12$ and is reasonably good. Higgs boson is predicted to be light, $m_H = 84_{-24}^{+32}$ GeV; at $\mu = m_Z$ the value of QCD coupling constant $\hat{\alpha}_s = 0.184(27)$.

3 Fits with extra generation

Introducing the fourth generation we assume that the mixing with three known generations is small. We fix the value of the sum of new quark masses $m_U + m_D = 600$ GeV to avoid Tevatron direct search bounds and the value of the charged lepton mass $m_E = 200$ GeV (well above LEP II bound) and look for a minimum of χ^2 for the fixed values of m_H varying neutral lepton mass and the difference of Up- and Down-quark masses. From the width of Z we know that neutral lepton should be heavy, $m_N > 50$ GeV. The results of fits for $m_H = 120$ GeV and $m_H = 1000$ GeV are shown on Figures 1 and 2.




Figure 1: *Exclusion plot on the plane $m_N, m_U - m_D$ for fixed values $m_H = 120$ GeV, $m_E = 200$ GeV, $m_U + m_D = 600$ GeV. χ^2 minimum shown by the star corresponds to $\chi^2/\text{d.o.f.} = 17.7/11$. The borders of the regions show domains allowed at the level $\Delta\chi^2 = 1, 4, 9, 16$ etc.*




Figure 2: *The same as Fig. 1 for $m_H = 1000$ GeV. χ^2 minimum shown by the star corresponds to $\chi^2/\text{d.o.f.} = 18.4/11$.*

We see that the quality of fits is almost the same as in the case of three generations.

4 S, T, U versus S_i, T_i, U_i

These variables are frequently used to determine if a given new physics model passed constraints of precision measurements.

Table

m_H		T_i	T	S_i	S	U_i	U
120	$m_U = 310, m_D = 290$	0.02	0.02	0.15	0.15	0	0
	$m_N = 120, m_E = 200$	0.11	0.11	-0.01	-0.01	0.02	0.01
1000	$m_U = 315, m_D = 285$	0.05	0.05	0.15	0.15	0	0
	$m_N = 53, m_E = 200$	0.27	0.36	-0.19	-0.13	0.16	0.10

Quark and lepton contributions to S, T, U and S_i, T_i, U_i at the points of χ^2 minimum in Figures 1 and 2. All masses are in GeV.

In the Table values S, T and U for the points of χ^2 minimum in Figures 1 and 2 are presented. Definitions of S, T and U from the review paper² are used. Also the values S_i, T_i and U_i where “ i ” means “improved” are presented. While used in² definitions of S, T and U correctly describe contributions of heavy new particles, $(m_{n.p.}/M_Z)^2 \gg 1$, quantities S_i, T_i and U_i can be used for $m_{n.p.} \lesssim M_Z$ as well (this improved set was introduced in¹ where it was designated as S', T' and U'). For the case of light higgs $m_H = 120$ GeV $S = S_i = 0.14, T = T_i = 0.13$ and both U_i and U are almost zero. In Fig. 10.4 of² one can see that both Standard Model point $S = T = 0$ and just described new physics point are on the border of the allowed 1σ domain for $m_H = 117$ GeV. This confirms the statement about equally good data description by SM and by the fourth family extension. For the case of heavy higgs $m_H = 1000$ GeV lepton contributions to S, T and U differs from that to S_i, T_i and U_i . Summing quark and lepton contributions we get: $S = 0.02, T = 0.41, U = 0.10$ and the (S, T) point is outside 1σ allowed domain for $m_H = 1000$ GeV from Fig. 10.4. However taken into account improved values $S_i = -0.04, T_i = 0.32, U_i = 0.16$ we see, that (S_i, T_i) point is at the border of 1σ allowed domain for $m_H = 1000$ GeV from Fig. 10.4 as it should be according to the analysis of Section 3. Similar statements can be found in³.

5 Higgs properties with the fourth generation

Loops with U - and D -quarks contribute to $gg \rightarrow H$ production amplitude. This helps CDF to exclude larger interval of H masses than in the case of 3 generations⁴. H decay branching ratios are modified as well⁵.

Additional heavy fermions generate negative loop correction to the Higgs boson potential which can make it unbounded from below. The allowed domains of fermion and Higgs boson masses from⁶ are shown on Fig. 3. They depend on the ultraviolet cut-off Λ at which new physics comes into play. One can see that in case the fourth generation exists we should have another type of relatively light ($\Lambda < 10^5$ TeV) new physics as well.

6 Conclusion

One extra quark-lepton generation is not excluded by electroweak precision data.

The quality of fit for one extra generation is the same as that for Standard Model for certain values of new particle masses.




Fig. 1. Allowed values of M_H and M_4 lie between two curves: a. solid for $\Lambda = 10^5$ GeV; b. thin dotted for $\Lambda = 10^{10}$ GeV; c. thick solid for $\Lambda = 10^{15}$ GeV; d. thick dotted for $\Lambda = 10^{19}$ GeV.

Figure 3: Bounds from the stability of Higgs potential and absence of the Landau pole.

In the case of the fourth generation the upper bound on higgs mass from Standard Model fit is removed. Higgs production cross-section in gluon fusion is enhanced while decay branching ratios are modified in the case of extra generations.

The fourth generation can be used to explain possible deviations from the Standard Model predictions for CP violation in B -meson decays^{7, 8, 9}. However presently I cannot find experimentally proved deviation which can not be attributed to the effects of strong interactions.

I am grateful to my coauthors Victor Novikov and Alexander Rozanov.

I am grateful to the organizers and in particular to Jean Tran Than Van, Blazenka Melic and Aleksander Khodjamirian for the possibility to participate in a very interesting and exciting conferences.

References

1. V.A. Novikov, A.N. Rozanov, M.I. Vysotsky, *hep-ph* 0904.4570, accepted to *Yad. Fiz.*
2. J. Erler, P. Langacker, *Review of Particle Physics*, in C. Amsler et al., *Phys. Lett. B* **667**, 1 (2008), chapter 10.7.
3. Graham D. Kribs, Tilman Plehn, Michael Spannowsky, Timothy M.P. Tait. *Phys. Rev. D* **76**, 075016 (2007).
4. M. Mulhearn, *Moriond QCD 2008*.
5. J.M. Frere, A.N. Rozanov, M.I. Vysotsky, *Phys. Atom. Nucl.* **69**, 355 (2006).
6. H.B. Nielsen, A.V. Novikov, V.A. Novikov, M.I. Vysotsky, *Phys. Lett. B* **374**, 127 (1996).
7. W.S. Hou, H. Li, S. Mishima, M. Nagashima, *Phys. Rev. Lett.* **98**, 131801 (2007).
8. A. Soni, A. Alok, A. Giri, R. Mohanta, S. Nandi, *hep-ph* 0807.1971.
9. M. Bobrowski, A. Lenz, J. Riedl and J. Rohrwild, *Phys. Rev. D* **79**, 113006 (2009).